

QED $\otimes$ QCD EXPONENTIATION: SHOWER/ME MATCHING AND IR-IMPROVED DGLAP THEORY AT THE LHC

B.F.L. WARD AND S.A. YOST

Department of Physics, Baylor University, Waco, TX, USA

We discuss the elements of QED $\otimes$ QCD exponentiation and its interplay with shower/ME matching and IR-improved DGLAP theory in precision LHC physics scenarios. Applications to single heavy gauge boson production at hadron colliders are illustrated.

BU-HEPP-06-07, Oct., 2006, *presented by B.F.L. Ward at ICHEP06*

Given the approaching turn-on of the LHC, the issue of precision predictions for the effects of multiple gluon and multiple photon radiative processes is a more immediate. A problem of some interest is the construction of an event-by-event precision theory of potential luminosity processes such as single heavy gauge boson production processes. Presuming the LHC luminosity experimental error to reach 2% [1], the attendant theoretical precision tag on processes such as single W,Z production should be $\frac{2}{3} - 1\%$ so that the theoretical error does not adversely affect the respective luminosity determination and the attendant precision LHC physics^a. Thus, we have developed [3] a theory of the simultaneous resummation of multiple gluon and multiple photon radiative effects, QED $\otimes$ QCD exponentiation, to realize systematically the needed higher order corrections on an event-by-event basis, in the presence of parton showers, to the desired accuracy.

We note that the new QED $\otimes$ QCD exponentiation theory is an exact resummation theory in the spirit of the original YFS exponentiation for QED and is an extension of the QCD exponentiation theory presented in Refs. [4]. One should compare the latter to the formal proof of exponentiation in non-Abelian gauge theories *in the eikonal ap-*

proximation as given in Ref. [5]: the results in Ref. [4] are in contrast *exact* but have an exponent that only contains the leading contribution of the exponent in Ref. [5]— if desired, all of the contributions of the latter exponent can be incorporated into the exponent Ref. [4] while maintaining exactness. Our precision requirement for the LHC luminosity processes requires exact $\mathcal{O}(\alpha_s^2, \alpha_s \alpha, \alpha^2)$ QED $\otimes$ QCD exponentiation in the presence of parton showers, on an event-by-event basis, realized by MC methods for realistic multiple gluon and multiple photon radiative effects. Development of such MC's is in progress.

We note that in Ref. [3] we have shown that for a process such as $q + \bar{q}' \rightarrow V + n(G) + n'(\gamma) + X \rightarrow \bar{\ell}\ell' + n(g) + n'(\gamma) + X$ we have the result

$$d\hat{\sigma}_{\text{exp}} = e^{\text{SUM}_{\text{IR}}(\text{QCD})} \sum_{n,m=0}^{\infty} \int \prod_{j_1=1}^n \frac{d^3 k_{j_1}}{k_{j_1}} \prod_{j_2=1}^m \frac{d^3 k'_{j_2}}{k'_{j_2}} \int \frac{d^4 y}{(2\pi)^4} e^{iy \cdot (p_1 + q_1 - p_2 - q_2 - \sum k_{j_1} - \sum k'_{j_2}) + D_{\text{QCD}}} \tilde{\beta}_{n,m}(k_1, \dots, k_n; k'_1, \dots, k'_m) \frac{d^3 p_2}{p_2^0} \frac{d^3 q_2}{q_2^0}, \quad (1)$$

where the new YFS [6, 7] residuals, defined in Ref. [3], $\tilde{\beta}_{n,m}(k_1, \dots, k_n; k'_1, \dots, k'_m)$, with n hard gluons and m hard photons, represent the successive application of the YFS expansion first for QCD and subsequently for QED. The functions $\text{SUM}_{\text{IR}}(\text{QCD})$, D_{QCD}

^aWe note for reference that the current precision tag for the luminosity at FNAL is at the $\sim 7\%$ level [2].

are given in Ref. [3]. The residuals $\tilde{\beta}_{n,m}(k_1, \dots, k_n; k'_1, \dots, k'_m)$ are free of all infrared singularities and the result in (1) is a representation that is exact and that can therefore be used to make contact with parton shower MC's without double counting or the unnecessary averaging of effects such as the gluon azimuthal angular distribution relative to its parent's momentum direction.

For our prototypical processes, we have the standard formula

$$d\sigma_{exp}(pp \rightarrow V + X \rightarrow \bar{\ell}\ell' + X') = \sum_{i,j} \int dx_i dx_j F_i(x_i) F_j(x_j) d\hat{\sigma}_{exp}(x_i x_j s), \quad (2)$$

in the notation of Ref. [3]. We will illustrate this formula with semi-analytical methods and the case of single Z production with the structure functions $\{F_i\}$ from Ref. [8] for definiteness. A MC realization will appear elsewhere [9].

As we have explained in Ref. [10], from the standpoint of using the result (1) with the structure functions $\{F_i\}$, we intend two possible approaches to the implied shower/ME matching [11]: one is based on p_T matching and one based on the shower-subtracted residuals $\hat{\tilde{\beta}}_{n,m}$. The respective shower can be that from PYTHIA [12], HERWIG [13], or the new shower algorithms in Ref. [14]. This combination of showers and exact ME's can be improved systematically to arbitrary precision order-by-order in (α_s, α) in the presence of exact phase space [9].

To illustrate the size of the interplay between QED and QCD in the threshold region, we compute, with and without the QED, the respective ratio $r_{exp} = \sigma_{exp}/\sigma_{Born}$, with the results

$$r_{exp} = \begin{cases} 1.1901 & , \text{QCED} \equiv \text{QCD} + \text{QED}, \text{ LHC} \\ 1.1872 & , \text{QCD}, \text{ LHC} \\ 1.1911 & , \text{QCED} \equiv \text{QCD} + \text{QED}, \text{ Tevatron} \\ 1.1879 & , \text{QCD}, \text{ Tevatron}. \end{cases} \quad (3)$$

Here, we have used from (1) the $\tilde{\beta}_{0,0}$ -level

result

$$\hat{\sigma}_{exp}(x_1 x_2 s) = \int_0^{v_{max}} dv \gamma_{QCED} v^{\gamma_{QCED}-1} F_{YFS}(\gamma_{QCED}) e^{\delta_{YFS}} \hat{\sigma}_{Born}((1-v)x_1 x_2 s) \quad (4)$$

where we intend the well-known results for the respective parton-level Born cross sections and the value of v_{max} implied by the experimental cuts under study. The value for the QED $\otimes$ QCD exponent is $\gamma_{QCED} = \left\{ 2Q_f^2 \frac{\alpha}{\pi} + 2C_F \frac{\alpha_s}{\pi} \right\} L_{nls}$ where $L_{nls} = \ln x_1 x_2 s / \mu^2$ when μ is the factorization scale. The functions $F_{YFS}(\gamma_{QCED})$ and $\delta_{YFS}(\gamma_{QCED})$ are well-known [7] as well:

$$F_{YFS}(\gamma_{QCED}) = \frac{e^{-\gamma_{QCED}\gamma_E}}{\Gamma(1 + \gamma_{QCED})}, \quad (5)$$

with $\delta_{YFS}(\gamma_{QCED}) = \frac{1}{4}\gamma_{QCED} + (Q_f^2 \frac{\alpha}{\pi} + C_F \frac{\alpha_s}{\pi})(2\zeta(2) - \frac{1}{2})$, where $\zeta(2)$ is Riemann's zeta function of argument 2, i.e., $\pi^2/6$, and γ_E is Euler's constant, i.e., 0.5772... .

We see that the size of the QED effects, $\sim 0.3\%$, is appreciable in a 1% precision tag budget and these effects are comparable to the QED effects found for the structure function evolution itself in Refs. [15–19]. We also see that our soft, QCD correction is entirely consistent with the exact results in Ref. [20–22] and that our QED effects are consistent with the exact $\mathcal{O}(\alpha)$ results in Refs. [23–25]. Such cross-checks are essential in establishing a rigorous precision tag on the theoretical predictions.

With an eye toward the 1% precision tag for the single heavy gauge boson production at the LHC, we have analyzed [26] the IR limit of the DGLAP [27] kernels themselves. The effects which we address are illustrated in Fig. 1. We apply the QCD exponentiation master formula from Ref. [4], the analog of (1) for just QCD, to the gluon emission transition that corresponds to $P_{qq}(z)$, i.e., to the squared amplitude for $q \rightarrow q(z) + G(1-z)$; as we show in Ref. [26], this allows us to get the following IR-improved DGLAP kernels

$$P_{qq}^{exp}(z) = C_F F_{YFS}(\gamma_q) e^{\frac{1}{2}\delta_q} \left\{ \frac{1+z^2}{1-z} (1-z)^{\gamma_q} - f_q(\gamma_q) \delta(1-z) \right\}, \quad (6)$$




Figure 1. In (a), we show the usual process $q \rightarrow q(1-z) + G(z)$; in (b), we show its multiple gluon improvement $q \rightarrow q(1-z) + G_1(\xi_1) + \dots + G_n(\xi_n)$, $z = \sum_j \xi_j$.

$$P_{Gq}(z)^{exp} = C_F F_{YFS}(\gamma_q) e^{\frac{1}{2}\delta_q} \frac{1 + (1-z)^2}{z} z^{\gamma_q}, \quad (7)$$

$$P_{GG}^{exp}(z) = 2C_G F_{YFS}(\gamma_G) e^{\frac{1}{2}\delta_G} \left\{ \frac{1-z}{z} z^{\gamma_G} + \frac{z}{1-z} (1-z)^{\gamma_G} + \frac{1}{2} (z^{1+\gamma_G} (1-z) + z(1-z)^{1+\gamma_G}) - f_G(\gamma_G) \delta(1-z) \right\}, \quad (8)$$

$$P_{qG}^{exp}(z) = F_{YFS}(\gamma_G) e^{\frac{1}{2}\delta_G} \frac{1}{2} \{ z^2 (1-z)^{\gamma_G} + (1-z)^2 z^{\gamma_G} \}, \quad (9)$$

where from the standard DGLAP methodology [26] we have $f_q(\gamma_q) = \frac{2}{\gamma_q} - \frac{2}{\gamma_q+1} + \frac{1}{\gamma_q+2}$, $\gamma_q = C_F \frac{\alpha_s}{\pi} t = \frac{4C_F}{\beta_0}$, $\delta_q = \frac{\gamma_q}{2} + \frac{\alpha_s C_F}{\pi} (\frac{\pi^2}{3} - \frac{1}{2})$, $\gamma_G = C_G \frac{\alpha_s}{\pi} t = \frac{4C_G}{\beta_0}$, $\delta_G = \frac{\gamma_G}{2} + \frac{\alpha_s C_G}{\pi} (\frac{\pi^2}{3} - \frac{1}{2})$, and

$$f_G(\gamma_G) = \frac{n_f}{C_G} \frac{1}{(1+\gamma_G)(2+\gamma_G)(3+\gamma_G)} + \frac{2}{\gamma_G(1+\gamma_G)(2+\gamma_G)} + \frac{1}{(1+\gamma_G)(2+\gamma_G)} + \frac{1}{2(3+\gamma_G)(4+\gamma_G)} + \frac{1}{(2+\gamma_G)(3+\gamma_G)(4+\gamma_G)}. \quad (10)$$

Here, $\beta_0 = 11 - \frac{2}{3}n_f$, where n_f is the number of active quark flavors. We show in Ref [26, 28] that these improved kernels change significantly the evolution of the moments of the structure functions. For example, the non-singlet anomalous dimension

A_n^{NS} becomes, from the exponentiated kernels,

$$A_n^{NS} = C_F F_{YFS}(\gamma_q) e^{\frac{1}{2}\delta_q} [B(n, \gamma_q) + B(n+2, \gamma_q) - f_q(\gamma_q)] \quad (11)$$

where $B(x, y)$ is the beta function given by $B(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x+y)$. Compare the usual result

$$A_n^{NS^o} \equiv C_F \left[-\frac{1}{2} + \frac{1}{n(n+1)} - 2 \sum_{j=2}^n \frac{1}{j} \right]. \quad (12)$$

The IR-improved n-th moment goes for large n to a multiple of $-f_q$, consistent with $\lim_{n \rightarrow \infty} z^{n-1} = 0$ for $0 \leq z < 1$; the usual result diverges as $-2C_F \ln n$. The two results differ for finite n as well: we get, for example, for $\alpha_s \cong .118$, $A_2^{NS} = C_F(-1.33)$, $C_F(-0.966)$ for (12) and (11), respectively. See Refs. [26, 28] for further discussion.

Contact with the exact 2-loop and 3-loop results in Refs. [29, 30] has also been made [26]: for the non-singlet case, we have, in the notation of Ref. [30], the exact 3-loop IR-improved result

$$P_{ns}^{+,exp}(z) = \left(\frac{\alpha_s}{4\pi} \right) 2P_{qq}^{exp}(z) + F_{YFS}(\gamma_q) e^{\frac{1}{2}\delta_q} \left[\left(\frac{\alpha_s}{4\pi} \right)^2 \{ (1-z)^{\gamma_q} \bar{P}_{ns}^{(1)+}(z) + \bar{B}_2 \delta(1-z) \} + \left(\frac{\alpha_s}{4\pi} \right)^3 \{ (1-z)^{\gamma_q} \bar{P}_{ns}^{(2)+}(z) + \bar{B}_3 \delta(1-z) \} \right] \quad (13)$$

where $P_{qq}^{exp}(z)$ is given in (9) and the resummed residuals $\bar{P}_{ns}^{(i)+}$, $i = 1, 2$ are related to the exact results [29, 30] for $P_{ns}^{(i)+}$, $i = 1, 2$, via

$$\bar{P}_{ns}^{(i)+}(z) = P_{ns}^{(i)+}(z) - B_{1+i} \delta(1-z) + \Delta_{ns}^{(i)+}(z), \quad (14)$$

where the B_i [29, 30] and the $\Delta_{ns}^{(i)+}(z)$ are given explicitly in Ref. [26]. The detailed phenomenological consequences of the fully exponentiated 2- and 3-loop DGLAP kernel set will appear elsewhere [9].

In summary, the methods illustrated and presented herein are all under investigation and implementation as we prepare for the era of 1% precision LHC theoretical predictions

on the necessary processes. Finally, one of us (B.F.L.W.) thanks Prof. W. Hollik for the support and kind hospitality of the MPI, Munich, while a part of this work was completed. We also thank Prof. S. Jadach for useful discussions. This work is partly supported by US DOE grant DE-FG02-05ER41399 and by NATO grant PST.CLG.980342.

References

1. M. Dittmar *et al.*, *Phys. Rev.* **D56**, 7284 (1997); M. Rijssenbeek, in *Proc. HCP2002*, ed. M. Erdmann,(Karlsruhe, 2002) p. 424; M. Dittmar, *ibid.*,p. 431.
2. S. Klimenko, in *Proc. HCP2002*, ed. M. Erdmann,(Karlsruhe, 2002) p. 413.
3. C. Glosser *et al.*,*Mod. Phys. Lett.* **A19**, 2113 (2004); B.F.L. Ward *et al.*, in *Proc. DPF 2004, Int. J. Mod. Phys.* **A20**, 3735 (2005); in *Proc. ICHEP04, vol. 1*, eds. H. Chen *et al.*,(World. Sci. Publ. Co., Singapore, 2005) p. 588.
4. B.F.L. Ward and S. Jadach, *Acta Phys.Polon.* **B33**, 1543 (2002); in *Proc. ICHEP2002*, ed. S. Bentvelsen *et al.*,(North Holland, Amsterdam, 2003) p. 275 ; *Mod. Phys. Lett.* **A14**, 491 (1999) ;D.B. DeLaney *et al.*, *Phys. Lett.* **B342**, 239 (1995) ; *Mod. Phys. Lett.* **A12**, 2425 (1997); *DPhys. Rev.* **D52**, 108 (1995); *ibid.***D66**, 019903(E) (2002).
5. J. Gatherall, *Phys. Lett.* **B133**(1983) 90.
6. D. R. Yennie *et al.*, *Ann. Phys.* **13**, 379 (1961); see also K. T. Mahanthappa, *Phys. Rev.* **126**, 329 (1962).
7. See also S. Jadach *et al.*,*Comput. Phys. Commun.* **102**, 229 (1997); *Phys. Rev.* **D55**, 1206 (1997); *Phys. Rev.* **D63**, 113009 (2001); *Comp. Phys. Commun.* **130**, 260 (2000); *ibid.***140**, 432, 475 (2001).
8. A.D. Martin *et al.*, *Phys. Rev.* **D51**, 4756 (1995).
9. S. Jadach *et al.*, to appear.
10. B.F.L. Ward and S.A. Yost, hep-ph/0509003, in *Proc. HERA-LHC Workshop*, CERN-2005-014, DESY-PROC-2005-01, eds. A. De Roeck and H. Jung,(CERN, Geneva, 2005) p. 304.
11. S. Frixione and B. Webber, *J. High Energy Phys.* **0206**, 029 (2002); S. Frixione *et al.*, *ibid.***0308**, 007 (2003).
12. T. Sjostrand *et al.*, hep-ph/0308153.
13. G. Corcella *et al.*, hep-ph/0210213.
14. S. Jadach and M. Skrzypek, *Acta Phys. Pol.* **B35**, 735 (2004); hep-ph/0504263, 0504205.
15. S. Haywood, P.R. Hobson, W. Hollik and Z. Kunszt, in *Proc. 1999 CERN Workshop on Standard Model Physics (and more) at the LHC, CERN-2000-004*, eds. G. Altarelli and M.L. Mangano,(CERN, Geneva, 2000) p. 122.
16. H. Spiesberger, *Phys. Rev.* **D52**, 4936 (1995).
17. W.J. Stirling, "Electroweak Effects in Parton Distribution Functions", talk presented at ESF Exploratory Workshop, *Electroweak Radiative Corrections to Hadronic Observables at TeV Energies* , Durham, Sept., 2003.
18. M. Roth and S. Weinzierl, *Phys. Lett.* **B590**, 190 (2004).
19. W. J. Stirling *et al.*, in *Proc. ICHEP04*, eds. H. Chen *et al.* (World Sci. Publ., Singapore, 2005) p. 527.
20. R. Hamberg *et al.*, *Nucl. Phys.* **B359**, 343 (1991).
21. W.L. van Neerven and E.B. Zijlstra, *Nucl. Phys.* **B382**, 11 (1992); *ibid.* **B680**, 513 (2004).
22. C. Anastasiou *et al.*,*Phys. Rev.* **D69**, 094008 (2004).
23. U. Baur *et al.*, *Phys. Rev.* **D57**, 199 (1998); *ibid.***D59**, 013002 (1998); *ibid.***D65** (2002) 033007.
24. S. Dittmaier and M. Kramer, *Phys. Rev.* **D65**, 073007 (2002).
25. Z. Zykunov,*Eur. Phys. J.* **C3**, 9 (2001).

26. B.F.L. Ward, hep-ph/0508140.
27. G. Altarelli and G. Parisi, *Nucl. Phys.* **B126**, 298 (1977); Yu. L. Dokshitzer, *Sov. Phys. JETP* **46**, 641 (1977); L. N. Lipatov, *Yad. Fiz.* **20**, 181 (1974); V. Gribov and L. Lipatov, *Sov. J. Nucl. Phys.* **15**, 438, 675 (1972); see also J.C. Collins and J. Qiu, *Phys. Rev.* **D39**, 1398 (1989) for alternative discussion of the lowest order DGLAP theory.
28. B.F.L. Ward, hep-ph/0602025, in *Proc. RADCOR05, Nucl. Phys. B Proc. Suppl.* **157**, 215 (2006).
29. E. G. Floratos *et al.*, *Nucl. Phys.* **B129**, 66 (1977); *ibid.* **B139**, 545 (1978); *ibid.* **B152**, 493 (1979); A. Gonzalez-Arroyo *et al.*, *Nucl. Phys.* **B153**, 161 (1979); A. Gonzalez-Arroyo and C. Lopez, *Nucl. Phys.* **B166**, 429 (1980); G. Curci *et al.*, *Nucl. Phys.* **B175**, 27 (1980); W. Furmanski and R. Petronzio, *Phys. Lett.* **B97**, 437 (1980); E. G. Floratos *et al.*, *Nucl. Phys.* **B192**, 417 (1981); R. Hamberg and W. Van Neerven, *Nucl. Phys.* **B379**, 143 (1992).
30. S. Moch *et al.*, *Nucl. Phys.* **B688**, 101 (2004); *ibid.* **B691**, 129 (2004), and references therein.